




Dermatopathology Slide Review Part 135

Paul K. Shitabata, M.D.
Dermatopathology Institute
Torrance, CA









What is the best diagnosis?


- A. Trichoepithelioma
- B. Trichoadenoma
- C. Trichilemmoma
- D. Trichoblastoma
- E. Trichofolliculoma


Dilated follicular cyst with multiple pilosebaceous units emptying
Into it




Cyst demarcated from surrounding dermis











What is the best diagnosis?


- A. Syringoma
- B. Trichilemmoma
- C. Clear cell basal cell carcinoma
- D. Trichoadenoma
- E. Eccrine poroma


Clear Cell Syringoma


Dermal proliferation of comma shaped ducts













Sneddon-Wilkinson Disease (Subcorneal Pustular Dermatosis)


Subcorneal pustule with neutrophils


No acantholysis


Minimal epidermal hyperplasia















What is the best diagnosis?

- A. Chondrosarcoma
- B. Calcifying aponeurotic fibroma
- C. Chondroid syringoma
- D. Chondrodermatitis nodularis helicis
- E. Chordoma


Chondroid Syringoma





Circumscribed
Dermal nodule

Chondroid matrix
With chondrocytes

Variable ductal
differentiation







Pertinent differential diagnostic considerations include all except:

- A. Acrodermatitis chronica atrophicans
- B. Morphea
- C. Radiation dermatitis
- D. Scleromyxedema
- E. Scleroderma

Answer-Scleromyxedema

- This is morphea with an element of morphea profunda. Histologically similar changes may be seen in scleroderma, radiation damage, and end-stage Lyme borreliosis.
- Scleromyxedema would exhibit dermal fibroplasia with mucinosis

“Square biopsy sign”

Dermal
sclerosis

Super-
Ficial
And deep
Perivascu-
Lar lympho-
Plasmocyti-
infiltrate

