The Histiocytoses

Paul K. Shitabata, M.D.

Director Dermatopathology

Harbor-UCLA Dermatology

Which disease may be associated with hematologic disorders?

- A. Diffuse normolipemic plane xanthoma
- B. Necrobiotic xanthogranuloma
- c. Verruciform xanthoma
- D. Xanthelasma
- E. Juvenile Xanthogranuloma

May have more than one answer

A. Diffuse normolipemic plane xanthoma B. Necrobiotic xanthogranuloma E. Juvenile Xanthogranuloma

Immunohistochemical markers of Langerhans cells include:

- A. CD1a
- **B**. CD4
- c. CD10
- D. CD35
- E. CD68

A-CD1a

Variants of Langerhans cell disease include:

- A. Hashimoto-Pritzker disease
- B. Letterer-Siwe disease
- c. Congenital self-healing reticulohistiocytosis
- D. Indeterminate cell histiocytosis
- E. Progressive nodular histiocytosis

- A. Hashimoto-Pritzker disease
- B. Letterer-Siwe disease
- C. Congenital self-healing reticulohistiocytosis

Incorrect associations include:

- A. Tendon xanthomas-Elevated HDL
- B. Necrobiotic xanthogranuloma-Paraneoplastic syndrome
- c. Verruciform xanthoma-HPV
- D. Diffuse normolipemic plane xanthoma-Castleman's disease

May have more than one answer

A. Tendon xanthomas-Elevated HDL C. Verruciform xanthoma-HPV

Associations with verruciform xanthoma include all except:

- A. Limb defects
- B. AIDS
- c. HSV
- D. Trauma
- E. Carcinoma in situ

- A. Limb defects
- B. AIDS
- D. Trauma
- E. Carcinoma in situ

Multicentric reticulohistiocytosis correct associations include:

- A. Histology similar to JXG
- B. CD1a positivity
- c. Destructive athritis
- D. Nail fold changes
- E. Hyperlipidemia

- A. Histology similar to JXG
- C. Destructive athritis
- D. Nail fold changes
- E. Hyperlipidemia