

# Noninfectious Vesiculobullous and Vesiculopustular Diseases

Paul K. Shitabata, M.D.  
Director of Dermatopathology  
Harbor-UCLA Dermatology

# What is the location of the blister cavity?

Disease	Intraepidermal	Subepidermal
Friction blister		
Second degree burn		
Suction blister		
Fogo selvagem		

# Answers

Disease	Intraepidermal	Subepidermal
Friction blister	+	
Second degree burn		Usually +
Suction blister		+
Fogo selvagem	+ Subcorneal	

# Histopathologic patterns associated with Grover's disease include all except:

- Bullous pemphigoid
- Hailey-Hailey disease
- Darier's disease
- Pemphigus vulgaris
- Pemphigus foliaceus

# Answer

- Bullous pemphigoid

# Pertinent histopathological differential diagnoses for graft versus host disease include all except:

- Eruption of lymphocyte recovery
- Erythema multiforme
- Lupus erythematosus
- Drug associated linear IgA disease
- Lichen planus

# Answer

- Drug associated linear IgA disease

# Location of inflammatory cells?

Disease	DEJ	Intra-epidermal	Stratum corneum
Miliaria rubra			
Miliaria profunda			
Miliaria crystallinaria			

# Answers


Disease	DEJ	Intra-epidermal	Stratum corneum
Miliaria rubra		+	
Miliaria profunda	+		
Miliaria crystallinaria			+

# Target Antigen in Pemphigus

Disease	Desmoglein 1	Desmoglein 3
Pemphigus vulgaris		
Pemphigus foliaceus		
Paraneoplastic pemphigus		
Drug-induced pemphigus		
IgA Pemphigus		

# Answers

Disease	Desmoglein 1	Desmoglein 3
Pemphigus vulgaris		+
Pemphigus foliaceus	+	
Paraneoplastic pemphigus	+	+
Drug-induced pemphigus	+	+
IgA Pemphigus	+	+



# Target Antigens in Subepidermal Bullous Diseases

Disease	BPAG1 (BP230)	BPAG2 (BP180)	S6beta4 integrin	Laminin 5	Type VII collagen
BP					
CP					
EBA					
PG					
LABD					

# Answers

Disease	BPAG1 (BP230)	BPAG2 (BP180)	S6beta4 integrin	Laminin 5	Type VII collagen
BP	+	+			
CP		+	+	+	(ocular) (anti-epi)
EBA					+
PG		+			
LABD		+			

